REMARKS

The claims have been amended to delete multiple dependencies, thereby reducing the overall filing cost, and to place them into proper U.S. claim format. Further, the claims submitted with this application to be prosecuted, and which are amended herein, reflect those claims as amended in the International Application and attached as annexes to the International Preliminary Examination Report (IPER). No new matter has been added.

The marked-up and clean substitute specifications submitted herewith include paragraph numbering, the introductory paragraph labelled "Cross-Reference to Related Applications," the final paragraph and other informal changes to place the application in proper U.S. format. No new matter has been added. Prosecution on the merits hereof is respectfully requested.

Respectfully submitted,

Date: Of the 17, 2005

Reg. No. 24,603

D. Peter Hochberg Co., L.P.A. 1940 E.6th Street – 6th Floor Cleveland, OH 44114-2294 (216) 771-3800 DPH/sfm

EXPRESS MAIL CERTIFICATION UNDER 37 CFR 1.10

"Express Mail" label number: EV713320020US Date of deposit 17 2005

I hereby certify that the paper(s) identified above, and any document(s) referred to as attached hereto, is being deposited with the United States Postal Service on the date indicated in an envelope as "Express Mail Post Office to Addressee" and is addressed: Mail Stop PCT, Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia, 22313-1450.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant

Stefan Bracht

Serial No.

:

Filing Date

Herewith

Examiner

Group Art Unit:

Title

MEDICAL ACTIVE SUBSTANCE PATCH WITH

REDUCED OPTICAL CONSPICUOUSNESS ON THE SKIN

(as amended herein)

Attorney File:

RO4101US (#90568)

Mail Stop PCT Commissioner for Patents P.O. Box 1450 Alexandria, Virginia, 22313-1450

MARKED-UP SUBSTITUTE SPECIFICATION

10/553708 JC09 Rec'd PCT/PTO 17 OCT 2005,

Medical Active Agent Patch Optically Less Visible on Skin Medical Active Substance Patch with Reduced Optical Conspicuousness on the Skin

CROSS-REFERENCE TO RELATED APPLICATIONS

[00001] This application is a National Stage application of International Application No. PCT/EP2004/003748, filed on April 8, 2004, which claims priority of German application number 103 17 692.6, filed on April 17, 2003.

BACKGROUND OF THE INVENTION

Field of the Invention

[00002] The present invention relates to medical active substance patches, particularly to transdermal therapeutic systems, comprising a monolayer or multilayer, active substance-containing matrix and a backing layer connected with [[said]] the matrix, [[said]] the active substance patches being distinguished by an improved optical appearance when being worn on the skin.

[00003] The invention further encompasses processes enabling the production of such active substance patches.

Description of the Prior Art

[00004] Many of the active substances or auxiliary agents suitable for use in the manufacture of active substance patches or TTSs show a tendency to discolour, for example to yellow. Such adverse changes may also occur during the application period. It is known, for instance, that nicotine patches gradually turn yellow.

[00005] The aforementioned changes are in most cases due to oxidative decomposition processes which progress upon contact with atmospheric oxygen and moisture especially during storage of the active substance patch or when it is being worn on the skin and which are promoted by action of light. Particularly affected by such processes are pharmaceutical active substances, antioxidants, various enhancers (i.e. substances promoting or accelerating transdermal active substance absorption), as well as oxidation-sensitive components of the pressure-sensitive adhesive that is present in the active substance patch, such as resin adhesives, for example.

[00006] The extent of active substance decomposition does not necessarily have an adverse effect on the pharmaceutical quality of the products, for instance if the resulting decomposition products amount to only fractions of a weight percent of the starting composition and if these decomposition products are toxicologically acceptable. Thus, discolouration often already affects a product cosmetically whereas the pharmaceutical quality is still unimpaired. Frequently, users or

patients especially in the case of medicaments associate such disadvantageous changes in the optical appearance of the active substance patches with defectiveness or deterioration, which causes a feeling of insecurity in those patients.

[00007] Often these changes are yellow, brown or red discolourations as typically appear in chemical decomposition. Even slight changes in colour may be interpreted by the users or patients as indicative of a deterioration of the quality of the medicament.

[00008] The problem of discolouration occurs particularly if the product, in fresh condition after manufacture, initially appears colourless or white to the human eye and the above-mentioned discolouration occurs only after a certain period of storage or while the patch is being worn on the skin. This is perceived by the users to be even more critical and potentially dangerous than a discolouration which has been there from the start and only becomes more intense during storage.

[00009] In the field of medical active substance patches, transparent and colourless patches represent the ideal case in respect of cosmetics since the user himself or other persons regard them as inconspicuous when applied to the user's skin. Users of medicinal patches generally prefer patches with such inconspicuous properties because they reduce the risk of other people becoming aware of the user's need for treatment and possibly finding out about his illness.

[000010] If for reasons of cosmetics a transparent design of an active substance patch does not make sense, for example because the ingredients are coloured or because of discolouration occurring during storage, it is possible to equip the patch with a non-transparent backing layer. During the application period, this backing layer then prevents [[that]] the colour or discolouration [[is]] from being optically perceived.

[000011] In the latter case, it is disadvantageous, however, that patches or TTSs equipped with a nontransparent backing layer are much more conspicuous at the site of application, that is, on the patient's skin, than transparent or colourless patches. A measure known from the state of the art and frequently applied consists in applying a skin-coloured lacquer to the nontransparent backing layer. This, however, leads to a further problem since it proves extremely difficult to find a skin tone that in equal measure suits a larger number of users of different skin colour tone and is cosmetically acceptable. Taking into consideration all of the skin types of the world population, it is entirely impossible to determine a unitary,

opaque skin colour tone that would be suitable as the colour tone for a non-transparent backing layer. This problem could be solved, it is true, by producing otherwise identical active substance patches having differently coloured backing layers that match the different skin colour tones of the world population, but this is out of the question because of the complex manufacturing and distribution logistics, and ultimately for reasons of cost.

SUMMARY OF THE INVENTION

[000012] The object of the present invention was therefore to provide active substance patches which despite colourations that are already existent or occurring over time ensure an optically inconspicuous appearance of the patch especially when the patch is located at the application site. The intention here is to preferably find a uniform solution which is suitable for the most different skin colour tones of the world population.

[000013] A further object of the invention was to indicate processes by means of which such active substance patches can be obtained.

[000014] These objects are achieved by means of medical active substance patches according to elaim 1 the present invention and by means of processes of production according to the present invention elaim 13, as well as by means of the embodiments described in the dependent claims.

[000015] Thus, the above-mentioned disadvantages do not occur or only occur in attenuated form in the medical active substance patches described in the introductory part of claim-1 according to the present invention if the active substance patch is transparent or at least translucent and if - in the state of having been applied to a person's skin – [[said]] the patch, in an area of the skin covered with the patch, has a lightness colour value L₁ which is not less than 50% and not more than 200% of a lightness colour value L₂, L₂ being the lightness value of the region of the skin of the same person which surrounds the applied patch, and if the same is true in respect of the skin of a second or any other person, provided that L₂, for all the persons mentioned, is in a range from 5° to 100°, especially in a range from 20° to 90°. The aforementioned differences between the lightness values L₁, L₂ can be determined by measurements in representative spot checks of people of the respective skin type.

[000016] The colour value of the lightness L, designated as "lightness colour value" is a colorimetric characteristic value which, in conjunction with other characteristics, is used in engineering for the non-ambiguous characterisation of

colours. The lightness colour value is indicated in degrees and can be determined by colour measuring instruments. The values of colour lightness indicated herein were determined by means of a "tristimulus colorimeter CP-320" of the firm of Techkon GmbH (DE-61462 Königstein).

[000017] Surprisingly, it emerged that active substance patches having the aforementioned features of the invention were of inconspicuous appearance at the place of application, i.e. on the skin, and that such active substance patches are optically inconspicuous on the most different skin colour types of the world population. For example, an active substance patch of the invention has an equally inconspicuous optical appearance when applied to the skin of a user of Caucasian, light skin colour or to the skin of a user of dark, Negroid skin colour. For this reason, according to a preferred one embodiment an active substance patch of the invention is characterized in that comprises the lightness colour value L₂ of the [[said]] first person, measured in the area of the skin not covered by the patch, is the lightness colour value of a person of light, Caucasian skin colour, and that the lightness colour value L₂ of the [[said]] second person is the lightness colour value of a person of dark, negroid skin colour, or vice versa.

[000018] The manufacture of active substance patches having the features of the present invention introductory part of claim 1 and the substances suitable for [[said]] the manufacture are in principle known to those skilled in the art.

Substances which may be used to produce the matrix layer(s) are, for instance, from the group of the polyacrylates, poly(meth)acrylates, adhesive resins, cellulose derivatives, polyisobutylenes, styrene-isoprene-styrene block copolymers, styrene-butadiene-styrene block copolymers, polydimethyl siloxane, ethylene vinyl acetate copolymers and vinyl acetate, optionally with addition of auxiliary substances known to the skilled artisan. At least one of the matrix layers contains an active substance, the term active substance referring, in particular, to a pharmaceutical active substance or a plurality of such substances.

[000019] The active substance patches of the invention, which are composed of a matrix and a superimposed backing layer, are substantially transparent or at least translucent (i.e. transmitting light but not transparent) and in any case not opaque. Thus, the backing layer is also substantially transparent or translucent.

[000020] Suitable as a backing layer are, first of all, polyesters, such as polyethylene terephthalate (PET) and polybutylene terephthalate, but also almost any other skin-compatible plastics, such as polyvinyl chloride, ethylene vinyl

acetate, vinyl acetate, polyethylene, polypropylene, cellulose derivatives and many others.

[000021] According to a preferred one embodiment, the active substance patches of the invention contain one or more substances from the group of the dyes and pigments in at least one of their layers. In combination with the transparent or translucent properties of the patch it is thereby achieved that a colouration of the matrix ingredient(s) which has been existing from the start or a discolouration of [[said]] the ingredients which has begun and intensifies only after the patch has been manufactured is optically masked. At the same time, the colour is thereby sufficiently adapted to the skin tone of the application site so that the patch will be inconspicuous on the most different skin colour types. Preferably the The substance(s) used for optical masking, which are selected from the group of the dyes and pigments, are contained in the matrix layer or in at least one of the matrix layers of a multilayer patch.

[000022] According to a further, particularly preferred embodiment, optical masking is achieved by providing the transparent or translucent backing layer with a content of at least one substance selected from the group of the dyes and pigments. This can be accomplished, in particular, by coating the backing layer of the patch on its outer side, that is, on the side averted from the skin, with a coating or a lacquer which contains at least one dye or/and at least one pigment. This variant has the additional advantage that the dye(s) or pigment(s) cannot come into contact with the active substance-containing matrix.

[000023] It may further be of advantage advantageous for both the matrix layer(s) and the backing layer to contain a dye or dyes and/or a pigment or pigments.

[000024] It has, surprisingly, emerged that it is not so much the adaptation of the dyes or pigments to the respective skin tone which is decisive, rather this effect is essentially determined by the concentration(s) of the dyes or/and pigments utilised. The optical conspicuousness of an active substance patch is substantially determined by the concentrations of the dyes and pigments contained therein. In addition, the layer thickness of the patch must be taken into consideration in this connection. In order for the patch to be inconspicuous to the eye of a beholder, certain concentrations of the dyes or/and pigments (inclusive of the coloured or discoloured ingredients, in particular active substances) must not be exceeded.

These concentrations can be determined by means of the disclosed conditions mentioned in claim 1.

[000025] At low concentrations of the coloured or discoloured ingredients comprised in the matrix, even such dyes or pigments can still be optically inconspicuous as clearly deviate from the colour tone of the underlying skin at the place of application. The same applies if the patch is of a small layer thickness. The low concentration and/or the small layer thickness results in a scope for the concentration or in corresponding possibilities of varying the layer thickness, thus fulfilling the requirements for optical masking of discolourations of ingredients of active substance patches by admixing dyes or/and pigments.

[000026] A further improvement of the optical appearance of active substance patches applied to the skin can be achieved, according to a particularly preferred an additional embodiment, by providing at least [[that]] the surface of the backing layer which is averted from the skin with reduced reflection properties. This can either be accomplished by means of physical methods or by applying an antireflection layer or antireflection coating. Such a layer or coating preferably contains an optical dulling agent or a combination of at least two dulling agents. This antireflection layer may at the same time contain a dye or dyes or/and a pigment or pigments to mask the ingredients of the patch, as described above.

[000027] In addition, it is possible by matting to eliminate or reduce [[that]] the cause of optical conspicuousness of an active substance patch which is due to light reflection. Such light reflection frequently occurs in active substance patches which are provided with a transparent backing layer of smooth surface structure. The reflection properties of these backing layer materials differ greatly from the reflection properties of human skin, which is why such plasters are visually very conspicuous on the skin.

[000028] The active substance patches of the present invention are particularly advantageous if at least one layer of the matrix comprises one or more coloured ingredient(s). This may, in particular, be a substance or substances which is/are colourless in its/their initial state and which has/have a tendency to discolour or which discolour during storage or during the application period. Particularly preferred are active substance patches which contain one or more pharmaceutically active substances as coloured ingredients or as ingredients which have a tendency to discolour, with particular preference such as for nicotine.

[000029] Preferably, the <u>The</u> active substance patches mentioned [[are]] <u>may</u> be transdermal therapeutic systems. These are distinguished by enabling a constant delivery of active substances via the skin for a determined period of time. The structure and manufacture of such systems are in principle known to those skilled in the art.

[000030] The <u>present</u> invention further encompasses processes for the production of the above-described active substance patches. These processes comprise the following steps:

- (a) producing a system comprising a mono- or multilayer active substance-containing matrix and a backing layer connected therewith, wherein the matrix is produced by using (a) matrix polymer(s), (an) active substance(s) and auxiliary agents, and wherein one or more substance(s) selected from the group of the dyes and pigments is/are admixed to the matrix or/and the backing layer;
- (b) producing at least one further system according to step (a), this system being different in terms of the concentration of the dyes or/and pigments, and/or in terms of the type of the dyes or/and pigments used;
- (c) producing surface sections or punched pieces from the systems obtained in steps (a) and (b);
- (d) producing or providing colour charts with lightness colour values L_2 in the range from 5° to 100°, particularly in the range from 20° to 90°,
- (e) applying or affixing the sections or systems obtained in step (c) to the colour charts mentioned in (d);
- (f) measuring the colour values of the lightness L_1 of the systems located on the colour charts and determining the difference between L_2 and L_1 in each particular case;
- (g) selecting those systems with a colour value of the lightness L_1 which is not less than 50% and not more than 200% of the lightness colour value L_2 .
- [000031] Through the teaching of the present invention it is made possible to produce active substance patches which despite containing coloured or discolouring ingredients are not easily perceivable to an observer and are optically inconspicuous when being worn on the skin, independently of whether the patch is attached to the skin of a light-skinned or dark-skinned person.

BRIEF DESCRIPTION OF THE DRAWINGS

[000032] Figure 1 is a bar graph depicting individual test patches Nos. 1-8 in the order of their visual inconspicuousness.

DETAILED DESCRIPTION OF THE INVENTION

[000033] The invention will be illustrated in greater detail by means of the following examples.

Examples

[000034] 1. Preparation of backing layers of different pigment concentrations

[000035] Coating compounds were prepared from ethyl cellulose and different portions of a pigment mixture (see Table 1) and these compounds were coated by means of a doctor knife to a PET film of 15 μ m thickness (weight per unit area 7-10 g/m²).

[000036] Table 1:

No.	Ethyl cellulose	Pigment mixture	
	[%-wt.]	[%-wt.]	
1	99.75	0.25	
2	99.5	0.5	
3	99.0	1.0	
4	98.0	2.0	
5	96.0	4.0	

[000037] Pigment mixture:

50.0%-wt. of Naturell BB Plv Pigment

50.0%-wt. of Naturell Pulver Pigment

(from Cosnaderm Chemische Rohstoffe GmbH, D-68526 Ladenburg)

[000038] Used as control examples were:

- (6) PET film, aluminised und nicotine-resistant (nontranslucent)
- (7) PET film, 15 μm, transparent
- (8) Scotchpak 1006

[000039] 2. Preparation of skin patches

[000040] Skin patches were produced using the backing layers prepared under

1. To this end, Durotak DUROTAK® 2052 (National Starch & Chemical B.V.) was spread at a weight per unit area of 80 g/m² and in each case covered with one of the backing layers mentioned under 1. Subsequently, individual patches, each patch of a size of 1 cm², were punched out.

[000041] 3. Preparing colour charts corresponding to the human skin colours

[000042] By [[means]] use of the software "PowerPoint" (Microsoft) and a colour printer (HP-C LaserJet 4500; Hewlett-Packard) eight colour charts were established representing the various skin colour tones of the world population.

[000043] The colour tones of the colour charts are characterized set forth in "PowerPoint" by the six parameters colour tone, red, green, blue, saturation, intensity as listed below, and can be reproduced by means of these parameters:

[000044] Table 2:

Colour chart	Colour	Red	Green	Blue	Saturation	Intensity
No.	tone					
A	16	255	215	191	255	223
В	21	50	25	0	255	25
С	21	80	40	0	255	40
D	21	255	236	217	255	236
E	21	197	137	77	130	137
F	21	72	36	0	255	36
G	21	117	78	39	128	78
Н	25	255	226	183	255	219

[000045] The colour charts Nos. A-H were measured using a "tristimulus colorimeter CP-320" of the firm of Techkon GmbH (DE-61462 Königstein). The values (in degrees) for the lightness L, the red-green axis a, and the yellow-blue axis b were determined. For each colour chart, 10 measurements were made and the mean values determined. The mean values are represented in the following Table 3.

[000046] Table 3:

Colour chart No.	L Value	a Value	b Value
	(L ₂)		
A	82.464	10.986	13.634
В	21.791	-3.203	8.877
С	25.776	5.905	14.758
D	88.086	4.945	9.572
E	50.596	10.893	36.304

F	25.811	3.747	12.968	
G	32.562	5.519	21.015	
Н	83.228	6.712	24.95	
Mean value*	51.289	5.688	17.758	

^{*} These are the respective mean values determined using the values of the 8 colour charts.

[000047] As can be seen, the colour value of the lightness L varies most, whereas the "a" value differs only slightly.

[000048] The range of the skin colours for which the principle of the present invention can be advantageously employed, according to the above-described "L,a,B" system particularly comprises the range of "5,8,60" up to "100,4,0".

[000049] 4. Determining the differences in lightness value

[000050] The punched skin patches described under 2. were adhered to the colour charts described under 3. Subsequently, the lightness colour values L_1 of the affixed patches were determined using the measuring method described under 3. From the measurement values L_1 obtained, the difference to the lightness value L_2 of the respective background (i.e. the colour chart) was determined in each case. The percentage differences between the lightness colour values L_1 of the patch types (Nos. 1 to 5 and controls Nos. 6 to 8) affixed to the colour charts A to H on the one hand and the lightness values L_2 of the respective colour charts A to H on the other hand are represented in Table 4.

[000051] It is evident therefrom that on all the colour charts the transparent PET film (7) shows the smallest deviations in respect of the lightness value (positive control).

[000052] Conversely, the largest deviations were found in the control examples (6) and (8).

[000053] 5. Visual Evaluation

[000054] Since it is known that the colour perception of humans can deviate from the colorimetrically determined data, a visual assessment of the test patches affixed to the colour charts A to H, inclusive of the comparison examples 6 to 8, was carried out by test subjects.

[000055] To this end, a certain number (e.g. 10) of each of the test patches (1 to 8) was affixed to the colour charts A to H. These colour charts were presented to a group of test subjects under standardized conditions (lighting, distance, time for

observing). The number of the patches that were not detected by the probands was used – after statistical evaluation of the data – as a measure for the optical inconspicuousness and thereby the effectiveness of the optical masking of a patch.

[000056] In Figure 1 the individual test patches Nos. 1-8 are represented in the form of a bar chart in the order of their visual inconspicuousness (vertical axis). Patch No. 1 and control patch No. 7 were not perceivable or hardly perceivable on most of the colour charts.

[000057]

Table 4

L A 2.733 B 13.735 C 3.236 D 5.244 E 1.435 F 12.254 G 1.388	PET film 15 pm, alum. A 1.42 B 273 C 215 D 7.89 E 60.3 F 215 G 150
H 5.022 Scotchpak 1006 (No. 8)	H 2.65 0.25 % Pigment (No. 1)
A 2.903 B 267.078 C 210.289 D 9.157 E 58.254 F 209.79 G 144.61 H 4.059	A 6.5 B 25.5 C 12.6 D 8.13 E 6.35 F 17.8 G 7.07 H 7.51
0.50 % Pigment (No. 2)	1.00 % Pigment (No. 3)
A 6.978 B 22.133 C 7.294 D 5.325 E 4.577 F 17.144 G 4.53 H 3.189	A 9.94 B 7.81 C 1.51 D 10.6 E 6.4 F 10.8 G 3.19 H 10.2
2.00 % Pigment (No. 4)	4.00 % Pigment (No. 5)
L A 14.387 B 1.647 C 4.058 D 8.295 E 33.528 F 1.794 G 2.709 H 13.731	L A 9.29 B 54.5 C 19.2 D 21.6 E 4.53 F 28.4 G 17.9 H 20

[000058] What has been described above are preferred aspects of the present invention. It is of course not possible to describe every conceivable combination of components or methodologies for purposes of describing the present invention, but one of ordinary skill in the art will recognize that many further combinations and permutations of the present invention are possible. Accordingly, the present invention is intended to embrace all such alterations, combinations, modifications, and variations that fall within the spirit and scope of the appended claims.